



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

and in the right angled triangles  $bBO$ ,  $bfO$ , the side  $BO =$  the side  $Of$ , since they are radii of the same circle, and the side  $bO$  common, therefore the other angles are equal, that is  $O\hat{b}B = O\hat{b}f$ ; also in the right triangles  $bfO$ ,  $bfC$ , the side  $Of =$  the side  $fC$  by construction and the side  $bf$  is common, therefore the angle  $O\hat{b}f =$  the angle  $C\hat{b}f$ . But  $O\hat{b}f$  was shown to be  $= O\hat{b}B$ ; therefore  $C\hat{b}f = O\hat{b}B$ , that is, the three angles are equal to each other. But the three angles  $C\hat{b}f$ ,  $f\hat{b}O$ ,  $O\hat{b}B$ , make up the angle  $C\hat{b}B = HOX$  the given angle.

Now in the right angled triangles  $bfC$  and  $OIC$  the angle  $bCO$  is common to the two triangles, therefore the remaining angles are equal, that is, the angle  $COI =$  the angle  $C\hat{b}f$ ; but  $C\hat{b}f$  has been shown to be equal to  $O\hat{b}B$ , therefore  $COI = O\hat{b}B$ ; but  $O\hat{b}B$  is one-third of  $C\hat{b}B$  or  $HOX$ , therefore  $HOC =$  one-third  $HOX$ .

WM. HILLHOUSE.

New Haven, Conn., Jan., 1878.

SOLUTION OF PROB. 405 BY PROF. C. A. VAN VELZER.—To fix the idea take the determinant of the fourth order

$$\begin{vmatrix} a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \\ a_4 & b_4 & c_4 & d_4 \end{vmatrix} = 0.$$

To the second column add  $l_1$  times the first,  $m_1$  times the third, and  $n_1$  times the fourth, where  $l_1, m_1, n_1$  are chosen to satisfy the equations

$$\begin{aligned} (l_1-1)a_1 + m_1c_1 + n_1d_1 + b_1 &= 0, \\ (l_1-1)a_2 + m_1c_2 + n_1d_2 + b_2 &= 0, \\ (l_1-1)a_3 + m_1c_3 + n_1d_3 + b_3 &= 0. \end{aligned}$$

These three equations are sufficient to determine the values of  $(l_1-1), m_1, n_1$ , but if to these we add a fourth

$$(l_1-1)a_4 + m_1c_4 + n_1d_4 + b_4 = 0$$

these *four* equations form a consistent set in  $(l_1-1), m_1, n_1$ , since the determinant of the coefficients (viz. the original determinant) vanishes.

We see that by this first transformation the determinant reduces to

$$\begin{vmatrix} a_1 & a_1 & c_1 & d_1 \\ a_2 & a_2 & c_2 & d_2 \\ a_3 & a_3 & c_3 & d_3 \\ a_4 & a_4 & c_4 & d_4 \end{vmatrix}$$

Now to the second row of this determinant add  $l_2$  times the first,  $m_2$  times the third and  $n_2$  times the fourth, where  $l_2, m_2, n_2$  satisfy the equ'n's

$$\begin{aligned}(l_2-1)a_1+m_2a_3+n_2a_4+a_2 &= 0, \\(l_2-1)c_1+m_2c_3+n_2c_4+c_2 &= 0, \\(l_2-1)d_1+m_2d_3+n_2d_4+d_2 &= 0.\end{aligned}$$

By this transformation the determinant is changed into

$$\begin{vmatrix} a_1 & a_1 & c_1 & d_1 \\ a_1 & a_1 & c_1 & d_1 \\ a_3 & a_3 & c_3 & d_3 \\ a_4 & a_4 & c_4 & d_4 \end{vmatrix}$$

a determinant in which the first two rows are identical and also the first two columns. The same process evidently applies to determinants of any order.

---

### NEW NOTATION FOR ANHARMONIC RATIOS.

---

BY PROF. WILLIAM WOOLSEY JOHNSON.

1. THE notation here proposed has for its object to express in a symmetrical manner the relation between the six distinct values of the anharmonic ratio of four points and the four modes of writing each which constitute the 24 arrangements of the four letters concerned.

2. The anharmonic ratio of the section of  $AB$  by  $PQ$  is the ratio

$$\frac{AP}{BP} : \frac{AQ}{BQ};$$

regarding  $A$  and  $B$  as fixed points of reference, the constituent ratios  $AP \div BP$  and  $AQ \div BQ$  may be called the *position ratios* of  $P$  and  $Q$ . A distinction of sign being made between  $AP$  and  $PA$ , each value of the position ratio determines the position of a point, negative position ratios corresponding to points between  $A$  and  $B$ ; while the position ratio of  $A$  is zero and that of  $B$  is infinity. The anharmonic ratio is the position ratio of  $P$  divided by that of  $Q$ , and regarding  $Q$  as a third fixed point of reference the value of the anharmonic ratio is a fixed multiple of the position ratio of  $P$ , and may be considered a coordinate determining the position of  $P$ , in such a manner that the coordinate of  $Q$  is unity.

3. Now let this anharmonic ratio be denoted by writing the four letters in a square form thus,

$$\begin{matrix} P & A \\ B & Q \end{matrix} = \frac{AP}{BP} : \frac{AQ}{BQ} = \frac{AP \cdot BQ}{BP \cdot AQ} = x, \quad (1)$$

in which it is to be remembered that the letters occupy the position given to their coordinates in the form

$$\begin{matrix} x & 0 \\ \infty & 1 \end{matrix}$$